

Measurement of the Ξ^-p Scattering Cross Sections at Low Energy

J.K. Ahn^{a,*} S. Aoki^b K.S.Chung^c M.S.Chung^d H.En'yo^e
 T.Fukuda^f H. Funahashi^g Y. Goto^e A. Higashi^h M. Ieiriⁱ
 T. Iijimaⁱ M. Inuma^{g,1} K. Imai^g Y. Itow^j J.M. Lee^{c,2}
 S. Makino^g A. Masaike^g Y. Matsuda^g Y. Matsuyama^h
 S. Mihara^g C. Nagoshi^h I. Nomura^k I.S. Park^d N. Saito^g
 M. Sekimotoⁱ Y.M. Shin^l K.S. Sim^d R. Susukita^e
 R. Takashima^m F. Takeutchiⁿ P. Tlustý^{h,3} S. Weibe^l
 S. Yokkaichi^g K. Yoshida^g M. Yoshida^e T. Yoshida^o
 S. Yamashita^p

^a*Department of Physics, Pusan National University, Pusan, 609-735, Korea*

^b*Faculty of Human Development, Kobe University, Kobe 657-8501, Japan*

^c*Department of Physics, Yonsei University, Seoul 120-749, Korea*

^d*Department of Physics, Korea University, Seoul 136-701, Korea*

^e*RIKEN, Institute for Physical and Chemical Research, Wako, 351-0198, Japan*

^f*Research Center for Physics and Mathematics, Osaka Electro-Communication University, Osaka, 572-8530, Japan*

^g*Department of Physics, Kyoto University, Kyoto 606-8502, Japan*

^h*Institute for Nuclear Study, University of Tokyo, Tokyo 188-8501, Japan*

ⁱ*IPNS, KEK, High Energy Accelerator Research Organization, Oho, 1-1, Tsukuba 305-0801, Japan*

^j*Institute for Cosmic Ray Research, University of Tokyo, Tokyo 188-8502, Japan*

^k*National Institute for Fusion Science, Nagoya 464, Japan*

^l*University of Saskatchewan, Saskatoon, Canada*

^m*Kyoto University of Education, Kyoto 612, Japan*

ⁿ*Kyoto Sangyo University, Kyoto 606, Japan*

^o*Department of Physics, Osaka City University, Osaka 558-8585, Japan*

^p*International Center for Elementary Particle Physics, University of Tokyo, Tokyo 113, Japan*

Abstract

In this paper we report cross-section measurements for Ξ^-p elastic and inelastic scatterings at low energy using a scintillating fiber active target. Upper limit on the total cross-section for the elastic scattering was found to be 24 mb at 90% confidence level, and the total cross section for the $\Xi^-p \rightarrow \Lambda\Lambda$ reaction was found to be $4.3^{+6.3}_{-2.7}$ mb. We compare the results with currently competing theoretical estimates.

Key words: Ξ^-p elastic scattering, Ξ^-p inelastic scattering

PACS: 21.80.+a, 25.80.Pw

Hyperon-nucleon interaction, particularly in $S = -2$ system, has attracted great attention since it may shed light on our understanding of the nature of SU(3) symmetry breaking in a baryon-baryon interaction. Its importance is more highlighted to account for a possible six-quark H -dibaryon, which has been as yet unobserved experimentally. In addition, the ΞN - $\Lambda\Lambda$ interaction accounts for the existence of a double hypernucleus, which is a gateway to a strange hadronic matter. For elementary Ξ^-p scattering, there is a scanty supply of bubble-chamber experimental data at higher energies above 1 GeV/ c [1,2,3,4].

Since no elementary cross sections below 1 GeV/ c are available to date, our attempt to measure the Ξ^-p elastic scattering and conversion processes is very important in testing competing theoretical-model predictions [5,6,7,8,9]. In particular, a measurement of the $\Xi^-p \rightarrow \Lambda\Lambda$ conversion cross section is crucial in assessing the stability of Ξ^- quasi-particle states in nuclei.

Our experiment was carried out using a separated 1.66 GeV/ c K^- beam at the KEK proton synchrotron. The momenta of outgoing particles were measured with a K^+ spectrometer of approximately 5 m length. The momentum resolution ($\Delta p/p$) was 0.5% (rms) at 1.2 GeV/ c , and the background ratio of π^+ to K^+ was estimated to be less than 0.2 %. The scintillating fiber (SCIFI) detector consisting of $0.5 \times 0.5 \text{ mm}^2$ plastic scintillating fibers acts as a primary interaction target and visual track detector for identifying ($K^-, K^+\Xi^-$) reactions, and secondary interactions of Ξ^- in the effective volume of $8 \times 8 \times 10 \text{ cm}^3$. The Ξ^- particles are produced by (K^-, K^+) reactions on free protons or bound protons in carbon nuclei. The SCIFI detector was read using two sets of the image-intensifier tubes (IIT), which were arranged orthogonally. The

* Corresponding author: jkahn@kaon.phys.pusan.ac.kr

¹ Present address : Department of Physics, Hiroshima University, Hiroshima 739-8526, Japan

² Present address : Korea Research Institute for Standards and Science, Daejeon 305-600, Korea

³ Present address : Nuclear Physics Institute of the Academy of Science, 250 68 Řeř, Czech Republic

track residual, defined as the distribution of the distance between the fitted straight track and each cluster, weighted by the brightness of each cluster, was about $290 \mu\text{m}$. The detailed description of the experimental methods appears in elsewhere [10,11,12].

The event configuration of Ξ^-p elastic scattering is characterized as two particle tracks from the Ξ^- interaction vertex and one of the two involving a kink. On the other hand, the $\Xi^-p \rightarrow \Lambda\Lambda$ reaction candidates were identified by observing two V topology tracks and one stopping track from the (K^-, K^+) primary vertex, as shown in Fig. 1.

Out of over six thousand $(K^-, K^+)\Xi^-$ events a total of two candidate events were selected as Ξ^-p elastic scattering. The Ξ^-p candidates were selected by requiring that the track length of a recoil proton be longer than 3 mm and that of a scattered Ξ^- be also longer than 3 mm. None of them has a visible Λ decay following the decay of a scattered Ξ^- : $\Xi^- \rightarrow \Lambda\pi^-$ and $\Lambda \rightarrow p\pi^-$. After imposing constraints on two-body $\Xi^-p \rightarrow \Xi^-p$ scattering, one of the two finally survived.

We also observed three candidate events involving two visible Λ decays from a stopping prong at the (K^-, K^+) vertex. The three events contain two Λ particles carrying unbalanced momenta in terms of two-body kinematics; the residual nucleus carries the unbalanced missing momentum. We therefore interpreted these events as due to the reaction $^{12}\text{C}(\Xi^-, \Lambda\Lambda)\text{X}$.




Fig. 1. Event sample for $\Xi^-p \rightarrow \Lambda\Lambda$ scattering process with sketch images

A great deal of attention was paid to estimate detection efficiencies. The detection efficiencies were calculated by a Monte-Carlo simulation based on

GEANT. The incident Ξ^- particles were generated to reproduce the measured Ξ^- momentum spectrum, and scattered off free protons. Energy loss in the SCIFI target medium was taken into account. The simulation involves reproducing the response of the SCIFI detector through hit digitization according to the individual fiber and fiber sheet configurations. The simulated Ξ^-p scattering events were scanned manually by eye in the same way as the analysis of real data. The scanning efficiency was studied as a function of track length and opening angle.

The acceptance was estimated by using a large number of the simulated events in terms of the Ξ^- momentum and c.m. angle of scattering. Proper cutoffs were imposed on the track length and opening angle according to the scanning results of the simulated events. The track length for all the charged particles was required to be longer than 3 mm and the Λ flight length longer than 5 mm. The opening angle between the scattered Ξ^- and the recoil proton should be larger than $\cos \theta_{\Xi p} = 0.96$.

We calculate upper limit on cross section for Ξ^-p elastic scattering based on the observation of one Ξ^-p elastic scattering event with an invisible Λ decay. If only incident and scattered Ξ^- particles are taken into account regarding Ξ^-p elastic scattering, the acceptance is flat over the range of c.m. scattering angles. However, as we discussed above, with the requirement of the visible Λ decay the acceptance is not flat in very forward and backward directions. We therefore obtain the differential cross section for the elastic scattering channel in the range of the Ξ^- scattering angle $-0.35 < \cos \theta_{\Xi p}^{\text{cm}} < 0.65$.

The differential cross sections for Ξ^-p scattering processes can be written as $2\pi \left(\frac{d\sigma}{d\Omega} \right) \Delta \cos \theta = \frac{N_{\Xi^-p} / \eta_{\Xi^-p}}{\rho \cdot L_{\Xi^-}^{\text{tot}}}$, where N_{Ξ^-p} denotes the number of Ξ^-p scattering events. The target density of free protons, $\rho = 4.88 \times 10^{22} \text{ cm}^{-3}$, can be rewritten as $\rho_0 N_A / W$, where ρ_0 is the density of the target material $(\text{CH})_n$ in g/cm^3 , N_A is the Avogadro's number, and the atomic weight, W , is given by 13 g/mol. The detection efficiency, η_{Ξ^-p} was obtained by a Monte-Carlo simulation, as described in the previous paragraph. The total track length of the incident Ξ^- particles is represented by the factor $L_{\Xi^-}^{\text{tot}} = \sum_{i=1}^{N_{\Xi^-}^{\text{tot}}} (L_i^{\Xi^-} - L_0) = \frac{N'_{\Xi^-}}{\eta_{\Xi^-}} \cdot (\sum L_i^{\Xi^-} / N'_{\Xi^-} - L_0)$, where $N_{\Xi^-}^{\text{tot}}$ denotes the total number of Ξ^- particles and $N'_{\Xi^-} = 3834$ indicates the analyzed number of Ξ^- particles. $L_i^{\Xi^-}$ stands for the track length of each Ξ^- . The effective target length of the analyzed event set was estimated to be 2.39 cm after correction due to the SCIFI target geometry. The factor η_{Ξ^-} denotes the ratio of the number of analyzed Ξ^- events to the total number of Ξ^- events, which was found to be 60.1%. The minimum cutoff length of Ξ^- track, L_0 , is 0.3 cm. The total track length of the incident Ξ^- particles, $L_{\Xi^-}^{\text{tot}}$, was then found to be $1.34 \times 10^4 \text{ cm}$.

Assuming Poisson statistics, we set upper limits on the cross sections at 90%




Fig. 2. Upper limits on the Ξp cross sections at 90% confidence level, indicated by arrows, are compared with theoretical estimates of RGM-F, RGM-H, FSS, Nijmegen-D, and SU_6 quark model. In the bottom panel a data point represents the result obtained from $\Xi^{-12}\text{C} \rightarrow \Lambda\Lambda$ reaction assuming the effective proton number of 3.5. Poisson statistical error is quoted only.

confidence level (2.44 for 0 event and 3.86 for 1 event with 50% background). The differential cross section for the Ξ^-p elastic scattering can be obtained in two ways: one is based on the observation of an event involving no Λ decay, the other is due to no observation of such an event that involves a visible Λ decay following the scattered Ξ^- decay. The detection efficiency factors η_{Ξ^-p} are 20.4% and 43.9%, respectively. For the calculation based on an observed event, we take 0.5 background events in the signal, simply because of taking the similar probability of quasifree elastic scattering from carbon nuclei into account. Upper limit on the differential cross sections for the elastic scattering channel in the range of $-0.35 < \cos \theta_{\Xi^-p}^{cm} < 0.65$ were then found to be 1.9 mb/sr (with Λ) and 2.1 mb/sr (without Λ) at 90% confidence level, respectively. The null result for the Ξ^-p elastic scattering with Λ decay provides more strict upper limit on the elastic scattering cross section. For simplicity, assuming an isotropic angular distribution, upper limit on the total cross section for $\Xi^-p \rightarrow \Xi^-p$ scattering process is 24 mb at 90% confidence level, as shown in Fig.2.

Regarding the $\Xi^-p \rightarrow \Lambda\Lambda$ reaction, we calculated the total cross-section with a null result. The detection efficiency was found to be 74% over the whole region of c.m. angles, and the branching ratio of two $\Lambda \rightarrow p\pi^-$ decays (41%) was taken into account. Upper limit on the total cross-section for $\Xi^-p \rightarrow \Lambda\Lambda$ process was obtained to be 12 mb at 90% confidence level. We also calculated the total cross section based on three observed events due to the reaction $^{12}\text{C}(\Xi^-, \Lambda\Lambda)X$. We

simply assumed a quasifree scattering process. The effective proton number for the $^{12}\text{C}(\Xi^-, \Lambda\Lambda)\text{X}$ reaction reflects the initial state interaction of Ξ^- through the $\Xi^-p \rightarrow \Xi^0n$ process and the sticking probability that either of two Λ particles or both stays inside a nucleus, thereby decaying weakly. In eikonal approximation, the effective nucleon number can be written as

$$N_{(\Xi^-, \Lambda\Lambda)} = (1 - \bar{S}_\Lambda)^2 \int_0^\infty 2\pi b db \int_{-\infty}^\infty dz \rho(\sqrt{b^2 + z^2}) \\ \times \exp \left[-\bar{\sigma}_{\Xi^-p \rightarrow \Xi^0n} \int_{-\infty}^z dz' \rho(\sqrt{b'^2 + z'^2}) \right],$$

where atomic mass $A = \int d\vec{r} \rho(\vec{r})$, where the nuclear radius r is given with the impact parameter b and the z coordinate. The nuclear density is of three parameter Fermi-distribution of $\rho(r) = (1 - wr^2/r_c^2)/(1 + \exp(r - r_c/a))$, where $w = 0.149$, $a = 0.5224$, and $r_c = 2.355$ fm. The Λ sticking probability was assumed to be $\bar{S}_\Lambda = 0.15$ [13]. The effective proton number $Z_{\text{eff}} = (Z/A) \times N_{\text{eff}}$ is then found to range from 3.0 to 4.1 with $\sigma_{\Xi^-p \rightarrow \Xi^0n} = 18$ mb (predicted by RGM models [5,6,7]) and 2 mb (by Nijmegen-D model [8]), respectively. The SU_6 quark model predicts $\sigma_{\Xi^-p \rightarrow \Xi^0n}$ to be about 15 mb at $p_\Xi = 0.5\text{GeV}/c$ [9]. Our previous experimental result (≈ 14 mb) for inelastic scattering channels involving both $\Xi^-p \rightarrow \Lambda\Lambda$ and $\Xi^-p \rightarrow \Xi^0n$ processes suggests that $\sigma_{\Xi^-p \rightarrow \Xi^0n}$ be of the order of 10 mb[14]. For $\sigma_{\Xi^-p \rightarrow \Xi^0n} = 10$ mb, the effective proton number was obtained to be 3.5. This value in turn gives the total cross-section for $\Xi^-p \rightarrow \Lambda\Lambda$ of $4.3_{-2.7}^{+6.3}$ mb (statistical error only).

We discuss possible systematic uncertainties in estimating the cross sections for Ξ^-p elastic scattering and conversion processes. The Ξ^-p cross sections measured in different ways are consistent with each other, which permit us to draw reliable conclusions on the low-energy Ξ^-p and $\Lambda\Lambda$ interactions. Special attention was paid to quantify the systematic uncertainties. The possible uncertainties are discussed as follows: (1) The total cross section measurement for the Ξ^-p elastic scattering was based on the assumption of an isotropic angular distribution. If one takes the distribution $d\sigma/d\cos\theta_{\Xi^-p}^{cm} \sim 1 + \cos^2\theta_{\Xi^-p}^{cm}$, then the total cross section would be 20% larger than the estimate based on an isotropic angular distribution. (2) Uncertainty of the detection efficiency was investigated by varying cutoffs on the minimum track length. It was found to be 5% change in the cross section results. (3) The SCIFI target is composed of almost equal amount of hydrogen and carbon, and also includes a tiny amount of other atomic components due to cladding and glue materials. This contribution was estimated to be of the order of 1% in the cross section results. It should be also noted that all the criteria for event selection were determined by scanning and analyzing a number of simulated events. All the systematic uncertainties are much smaller than limited significance of our statistics.

The total cross section for the $\Xi^-p \rightarrow \Lambda\Lambda$ conversion process sets tight limit on theoretical estimates as shown in Fig. 2. The upper limit on the cross section, 12 mb at 90% confidence level, rules out the RGM-F model prediction [5]. The measured cross section based on quasifree assumption is $4.3_{-2.7}^{+6.3}$ mb, which supports rather the prediction of the model $D(r_c = 0.5 \text{ fm})$ [8] of Nijmegen group and the SU_6 quark model [9]. Note that the quasifree scattering cross section depends on the estimate of the effective number of protons involved in the $^{12}\text{C}(\Xi^-, \Lambda\Lambda)X$ reaction. The above estimate was based on the effective proton number of 3.5, as discussed earlier. Due to this uncertainty of the effective number of protons, the models FSS [6] and RGM-H [7] cannot be ruled out.




Fig. 3. Cross sections for the $\Xi^-p \rightarrow \Lambda\Lambda$ reaction with respect to different reaction channels: (from left) $\Xi^-^{12}\text{C} \rightarrow \Lambda\Lambda X$ with three observed events, $\Xi^-p \rightarrow \Lambda\Lambda$ with a null result, $^{12}\text{C}(K^-, K^+)\Xi^-X$, and $^{12}\text{C}(K^-, K^+\Lambda\Lambda)X$ with intra-nuclear cascade model prediction for intermediate Ξ^- and meson-induced processes.

We discuss consistency of $\Xi^-p \rightarrow \Lambda\Lambda$ cross sections measured using four different reactions; $\Xi^-p \rightarrow \Lambda\Lambda$, $\Xi^-^{12}\text{C} \rightarrow \Lambda\Lambda X$, $K^-^{12}\text{C} \rightarrow K^+\Lambda\Lambda X$, and $K^-^{12}\text{C} \rightarrow K^+X$ (no Ξ^-). Among them the direct measurement of elementary $\Xi^-p \rightarrow \Lambda\Lambda$ process should be emphasized since it is free from theoretical assumptions. Despite the measurement yielded no observed event, it sets the stringent upper limit on the cross section to be 12 mb at 90% confidence level. Secondly, quasifree $\Xi^-^{12}\text{C} \rightarrow \Lambda\Lambda X$ reaction events involving two momentum-unbalanced Λ particles permit us to obtain the cross section of $4.3_{-2.7}^{+6.3}$ mb. The errors quoted indicate a 90% confidence level interval for Poisson signals. This result basically contains an assumption of the effective proton number being 3.5 in the eikonal approximation. The uncertainty in estimating the effective proton number is indicated as dashed-dotted arrow lines in the same column of Fig. 3 (3.8 mb for $Z_{\text{eff}} = 4$ and 7.5 mb for $Z_{\text{eff}} = 2$).

We also estimated the total cross section from the reactions, $K^-^{12}\text{C} \rightarrow K^+\Lambda\Lambda X$ and $K^-^{12}\text{C} \rightarrow K^+X$ without Ξ^- production. These processes however compete with intermediate meson-induced two-step processes [14]. Both cases could

also involve rather sizable contribution from $\Xi^-p \rightarrow \Lambda\Sigma^0$ due to Fermi motion of nucleons in ^{12}C above 0.57 GeV/ c . The ratio of the observed number of the Ξ^- particles to that of the $^{12}\text{C}(K^-, K^+)$ reaction events would provide upper limits on the inelastic Ξ^-N cross section. The ratio was found to be $80.1 \pm 2.4_{-1.3}^{+0.6}\%$ [12]. The total cross section for Ξ^-p inelastic scattering was then deduced from the Ξ^- absorption probability for ^{12}C , which is approximately 14 mb assuming an isotropic angular distribution. A recent measurement of quasifree $p(K^-, K^+)\Xi^-$ reaction in emulsion plate yields $12.7_{-3.1}^{+3.5}$ mb in the momentum range 0.4–0.6 GeV/ c [15], which is well consistent with our result. On the other hand, the analysis of the $\Lambda\Lambda$ invariant mass spectrum suggests that the $\Xi^-p \rightarrow \Lambda\Lambda$ cross section is of the order of several mb at $p_{\Xi^-} \sim 0.55$ GeV/ c [16].

In nuclear matter the Ξ^- decay width $\Gamma_{\Xi^-} \simeq (v\sigma)_{\Xi^-p \rightarrow \Lambda\Lambda} \cdot (\rho^0/2)$ [17] arrives at ~ 3 MeV, based on the result of $4.3_{-2.7}^{+6.3}$ mb. The term $\rho^0/2 \sim 0.08 \text{ fm}^{-3}$ denotes proton density and v is a Ξ^-p relative velocity.

In summary the KEK-PS measurement provides the first experimental data on low-energy Ξ^-p elastic and conversion cross sections in the range from 0.2 GeV/ c to 0.8 GeV/ c , which are less than 24 mb at 90% confidence level and of the order of several mb, respectively. The four independent and consistent measurements of $\Xi^-p \rightarrow \Lambda\Lambda$ cross section rule out a theoretical estimate based on the RGM-F model calculation, and favor the estimates predicted by Nijmegen model-D($r_c = 0.5$ fm) and SU_6 quark model. In nuclear matter, the width for a Ξ^- single-particle state is estimated to be $\Gamma_{\Xi^-} \approx 3$ MeV.

We would like to thank the staffs of the KEK proton synchrotron for their continuous support. We also thank Prof. Y. Fujiwara, Dr. C. Nakamoto and Dr. Y. Nara for valuable discussions and for providing us with theoretical calculation results. This work was supported in part by a Grant-in-Aid for Scientific Research (08239103), the Ministry of Science, Culture and Education, Japan. The work of J.K. Ahn was supported by Korea Research Foundation grant (KRF-2003-015-C00130).

References

- [1] G.R. Charlton, Phys. Lett. **B32**, 720 (1970).
- [2] R.D.A. Dalmeijer *et al.*, Nouvo Cimento Letters **4**, 373 (1970).
- [3] R.A. Muller, Phys. Lett. **B38**, 123 (1972).
- [4] J.M. Hauptman *et al.*, Nucl. Phys. **B125**, 29 (1977).
- [5] Y. Fujiwara, C. Nakamoto and Y. Suzuki, Prog. Theor. Phys. **94**, 353 (1995).

- [6] Y. Fujiwara, C. Nakamoto and Y. Suzuki, Phys. Rev. Lett. **76**, 2242 (1996).
- [7] Y. Fujiwara, C. Nakamoto and Y. Suzuki, Phys. Rev. **C54**, 2180 (1996).
- [8] C. Nakamoto, private communication ; C. Nakamoto, Y. Fujiwara and Y. Suzuki, Nucl. Phys. **A639**, 51c (1998); Y. Nara, private communication.
- [9] Y. Fujiwara, M. Kohno, C. Nakamoto and Y. Suzuki, Phys. Rev. **C64**, 054001 (2001).
- [10] J.K. Ahn *et al.*, Phys. Rev. **C62**, 55201 (2000).
- [11] J.K. Ahn, Ph.D thesis, Kyoto University *unpublished* (2000); Y. Itow, Ph.D thesis, Kyoto University *unpublished* (1994); S. Yamashita, Ph.D thesis, Kyoto University *unpublished* (1994).
- [12] M.S. Chung, Ph.D thesis, Korea University, *unpublished* (1995).
- [13] Y. Yamamoto, T. Motoba, T. Fukuda, M. Takahashi and K. Ikeda, Prog. Theor. Phys. Suppl. **117**, 281 (1994).
- [14] J.K. Ahn *et al.*, Nucl. Phys. **A625**, 231 (1997).
- [15] S. Aoki *et al.*, Nucl. Phys. **A644**, 365 (1998).
- [16] J.K. Ahn *et al.*, Phys. Lett. **B444**, 267 (1998).
- [17] C.B. Dover, A. Gal, and D.J. Millener, Nucl. Phys. **A572**, 85 (1994).